

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this

1. AGENCY USE ONLY (Leave blank)

2. REPORT DATE
27 April 1999

3. REPORT TYPE AND DATES COVERED
Final Report 1 Oct. 1995 - 30 Sept 98

4. TITLE AND SUBTITLE

Bioacoustics of Monterey Bay Pinnipeds:
Auditory fatigue and masking

5. FUNDING NUMBERS

N00014-95-1-0936

6. AUTHOR(S)

Dr. Ronald J. Schusterman

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Long Marine Laboratory
Institute of Marine Sciences
University of California at Santa Cruz
100 Shaffer Road
Santa Cruz, CA 95060

8. PERFORMING ORGANIZATION
REPORT NUMBER

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217-5000

10. SPONSORING / MONITORING
AGENCY REPORT NUMBER

11. SUPPLEMENTARY NOTES

12a. DISTRIBUTION / AVAILABILITY STATEMENT

Distribution Unlimited

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

Auditory thresholds were obtained for one northern elephant seal (*Mirounga angustirostris*), one harbor seal (*Phoca vitulina*) and two California sea lions (*Zalophus californianus*) under unmasked conditions, masked conditions, and following exposure to noise. Thresholds were determined using a 50% correct detection criterion in a signal detection task using trained behavioral responses to brief pure tones. Results indicate that 1) *Mirounga* is most sensitive to low frequency sounds in water and is relatively hard of hearing in air; 2) *Phoca* has comparable auditory sensitivity to sounds of low frequency in air and under water; 3) *Zalophus* is more sensitive to aerial sounds than underwater sounds; 4) *Zalophus* is susceptible to age related hearing loss, especially at higher frequencies; 5) all three species, and *Mirounga* in particular, are quite good at detecting low frequency signals from noise; and 6) moderately loud underwater sounds have the potential to induce short-term residual hearing loss in all species tested.

4. SUBJECT TERMS

pinniped, psychophysics, hearing, noise

15. NUMBER OF PAGES

4

16. PRICE CODE

17. SECURITY CLASSIFICATION
OF REPORT
unclassified

18. SECURITY CLASSIFICATION
OF THIS PAGE
unclassified

19. SECURITY CLASSIFICATION
OF ABSTRACT
unclassified

20. LIMITATION OF ABSTRACT

UL

19990505 121

FINAL REPORT

GRANT #: N00014-95-10936

PRINCIPAL INVESTIGATOR: Dr. Ronald J. Schusterman

INSTITUTION: University of California, Santa Cruz

GRANT TITLE: Bioacoustics of Monterey Bay pinnipeds: auditory fatigue and masking

AWARD PERIOD: 1 October 1995 - 30 September 1998

OBJECTIVE: To investigate the aerial and underwater hearing sensitivity of three pinniped species (*Zalophus californianus*, *Phoca vitulina*, *Mirounga angustirostris*) and to determine the effects of noise on hearing.

APPROACH: Auditory thresholds are determined for captive, trained animals using behavioral psychophysical techniques. Thresholds are obtained for three conditions: 1) unmasked (absolute sensitivity); 2) masked (in the presence of octave band noise); and 3) following exposure to noise (20 to 22 minutes of octave-band-noise centered at test frequency).

ACCOMPLISHMENTS: We established a series of low frequency (75 Hz to 6400 Hz) sound detection thresholds in air and underwater for *Zalophus*, *Phoca*, and *Mirounga*. Additionally, we obtained complete aerial and underwater audiograms for *Mirounga*, as well as complete underwater audiograms for a middle-aged *Zalophus* (12-year-old female) and an old *Zalophus* (23-year-old female). These data include the first hearing thresholds of any kind obtained for the deep diving *Mirounga*, and show that *Mirounga* hears extremely well at low frequencies (below 1000 Hz) in water but is relatively hard of hearing in air. Age related losses in hearing sensitivity were documented for *Zalophus* by comparing the two animals tested at mid-range (6000-8000 Hz) to relatively high (16,000-28,000 Hz) frequencies. The hearing deficit in the older animal at the mid-range frequencies was approximately 20dB, and the deficit at the high-range was between 40 and 60dB. Critical ratios, i.e., the amount by which masked thresholds exceed masking noise spectrum levels, were obtained at low frequencies (100 Hz to 2500 Hz) under nearly identical conditions for the three species. These data indicate that pinnipeds in general, and *Mirounga* in particular, are very good at differentiating signals from noise. In the first systematic experiments on underwater temporary threshold shift (TTS) in pinnipeds, hearing thresholds were obtained just prior to noise exposure, immediately following noise exposure, and within 24 hours of noise exposure. Test frequencies ranged from 100 Hz to 2000 Hz, and the octave-band noise exposure levels were approximately 60-75 dB SL (sensation level at center frequency). Each animal was trained to dive into a noise field and remain stationed underwater during a noise exposure period that lasted a total of 20-22 minutes. All subjects showed reliable shifts following noise exposure that averaged about 5 dB, and thresholds recovered to baseline levels within 24 hours. Control sessions, in which the subjects completed a simulated noise exposure, produced shifts that were reliably smaller than those produced following noise exposure. In the field, aerial and underwater

ambient noise levels were recorded at a harbor seal rookery and a northern elephant seal rookery, and acoustic signals were obtained from harbor seal pups, and northern elephant seal mothers and pups.

CONCLUSIONS: With respect to the absolute aerial and underwater hearing sensitivities of the species tested, we determined *Mirounga* to be the most "water-adapted", *Phoca* to be the most "amphibious-adapted", and *Zalophus* to be the most "air-adapted". Furthermore, the data show that elephant seals are most sensitive to sounds of low frequency under water. The critical ratios derived from the masking work for all three pinniped species are similar to those obtained for other marine mammals, and are generally lower than critical ratios obtained for terrestrial mammals. In particular, we are struck by the finding that our single northern elephant seal exhibited the lowest critical ratio of any mammal tested thus far at low frequencies. Reliable TTSS obtained for all three species under water indicate that moderately loud underwater sounds have the potential to cause short-term residual pinniped hearing impairment.

SIGNIFICANCE: These data facilitate prediction of auditory damage to pinnipeds resulting from exposure to noise. All three species are at risk to anthropogenic underwater noise at mid-range frequencies; however, *Mirounga* is at particular risk to low frequency underwater noise (below 1 kHz). With respect to airborne sound, *Zalophus* is particularly at risk to a range of frequencies (4-28 kHz), followed by *Phoca*. *Mirounga* is at little risk to aerial anthropogenic noise, because this species is generally hard of hearing throughout its entire aerial hearing range. The finding of age related hearing loss in *Zalophus* parallels findings in human subjects, in that sensitivity to higher frequencies is lost to a much greater extent than sensitivity to the lower frequencies. The masking data suggest that pinnipeds are efficient at extracting signals from ambient noise, even through they lack sonar. Field studies of ambient and biological noise is consistent with our lab work showing that *Mirounga* is relatively hard of hearing in air compared to *Phoca*.

PATENT INFORMATION: N/A

AWARD INFORMATION:

Ronald J. Schusterman, Fellow, Acoustical Society of America, 1998.

Ronald J. Schusterman, University of California at Santa Cruz, Department of Defense AASERT award for graduate student training, 1997-2000.

Ronald J. Schusterman, Fellow, Animal Behavior Society, 1997.

Brandon Southall, University of California at Santa Cruz, Outstanding Graduate Student in Ocean Sciences, 1997-1998.

PUBLICATIONS AND ABSTRACTS (for total period of grant):

1. Schusterman, R.J. and Kastak, D. (1995) There is no substitute for an experimental analysis of marine mammal cognition. *Marine Mammal Science*, 11, 263-267.

2. Hanggi, E.B. and Schusterman, R.J. (1995) Conditional discrimination learning in a male harbor seal (*Phoca vitulina*). In: Sensory Systems of Aquatic Mammals, R.A. Kastelein, J.A. Thomas and P.E. Nachtigall (Eds), De Spil Publishers, Woerden, Netherlands.
3. Kastak, D. and Schusterman, R.J. (1995) Aerial and underwater hearing thresholds for 100Hz pure tones in two species of pinnipeds. In: Sensory Systems of Aquatic Mammals, R.A. Kastelein, J.A. Thomas and P.E. Nachtigall (Eds), De Spil Publishers, Woerden, Netherlands.
4. Schusterman, R.J., Kastak, D. and Reichmuth, C. (1995) Equivalence class formation and cross modal transfer: testing marine mammals. In: Sensory Systems of Aquatic Mammals, R.A. Kastelein, J.A. Thomas and P.E. Nachtigall (Eds), De Spil Publishers, Woerden, Netherlands.
5. Handler, K.W., Belton, K., and Schusterman, R.J. (1995) Seasonal patterns of aerial aquatic displays and underwater vocalizations in Pacific harbor seals (*Phoca vitulina richardsi*). 11th Biennial conference on the Biology of Marine Mammals, Orlando, FL. (Abstract)
6. Kastak, D. and Schusterman, R.J. (1995) Comparative aspects of low frequency hearing in the northern elephant seal. 11th Biennial conference on the Biology of Marine Mammals, Orlando, FL. (Abstract)
7. Levenson, D.H. and Schusterman, R.J. (1995) Visual sensitivity relates to foraging environment in the northern elephant seal, California sea lion, and harbor seal: evidence from pupillometry. 11th Biennial conference on the Biology of Marine Mammals, Orlando, FL. (Abstract)
8. Reichmuth, C.J., Schusterman, R.J., and Kastak, D. (1995) Functional class formation as a cognitive mechanism in California sea lions. 11th Biennial conference on the Biology of Marine Mammals, Orlando, FL. (Abstract)
9. Kastak, D. and Schusterman, R.J. (1996) Temporary threshold shift in a harbor seal (*Phoca vitulina*). Journal of the Acoustical Society of America, 100:1905-1908.
10. Kastak, D. (1996) Comparative aspects of hearing in pinnipeds. Ph.D. Thesis, University of California at Santa Cruz.
11. Kastak, D. and Schusterman, R.J. (1996) Pinniped acoustics: adaptations for aerial and underwater hearing. Animal Behavior Society Meetings, Flagstaff, AZ (Abstract).
12. Schusterman, R.J. and Kastak, D. (1996) Pinniped acoustics: habituation and sensitization to anthropogenic signals. Animal Behavior Society Meetings, Flagstaff, AZ (Abstract)
13. Handler, K.W. and Schusterman, R.J. (1996) Inter-individual variation in display behavior of Pacific harbor seals at Point Lobos State Reserve, CA. Animal Behavior Society Meetings, Flagstaff, AZ (Abstract).
14. Reichmuth, C.J., Schusterman, R.J. and Kastak, D. (1996) Functional class formation as a cognitive mechanism in California sea lions. Animal Behavior Society Meetings, Flagstaff, AZ (Abstract).
15. Levenson, D.H., and Schusterman, R.J. (1996) Visual sensitivity relates to foraging behavior in pinnipeds: pupillometric evidence. Animal Behavior Society Meetings, Flagstaff, AZ (Abstract).

16. Schusterman, R.J., Kastak, D., and Reichmuth, C.J. (1997) What's in a name? Stimulus equivalence by any other name would smell as sweet. *Journal of the Experimental Analysis of Behavior*, 68, 252-258.
17. Reichmuth, C.J. (1997) Functional classes and equivalence relations in California sea lions. M.Sc. Thesis, University of California at Santa Cruz.
18. Levenson, D.H. and Schusterman, R.J. (1997) Pupillometry in seals and sea lions: ecological implications. *Canadian Journal of Zoology*, 75, 2050-2057.
19. Schusterman, R.J. (1997) Auditory sensitivity of a northern elephant seal to airborne and underwater sounds. 25th International Ethological Congress, Vienna, Austria. (Abstract)
20. Schusterman, R.J. (1997) Stimulus equivalence and animal social cognition. Animal Cognition Conference, Budapest, Hungary. (Abstract)
21. Kastak, D. and Schusterman, R.J. (1997) Aerial and underwater hearing sensitivity of a northern elephant seal (*Mirounga angustirostris*). 134th Meeting of the Acoustical Society of America, San Diego, CA. (Abstract)
22. Schusterman, R.J. and Kastak, D. (1998) Functional equivalence in a California sea lion: relevance to social and communicative interactions. *Animal Behavior*, 55:1087-1095.
23. Kastak, D. and Schusterman, R.J. (1998) Low frequency amphibious hearing in pinnipeds: methods, measurements, noise, and ecology. *Journal of the Acoustical Society of America*, 103, 2216-2228.
24. Kastak, D. and Schusterman, R.J. (1998) Pinniped underwater hearing: temporary threshold shifts in a harbor seal. *Animal Behavior Society Meetings*, Baltimore, MD. (Abstract)
25. Levenson, D.H. and Schusterman, R.J. (1998) Dark adaptation and visual sensitivity in seals and sea lions. *Animal Behavior Society Meetings*, Baltimore, MD. (Abstract)
26. Schusterman, R.J. (1998) New developments in marine mammal cognition and learning. The World Marine Mammal Science Conference, Monaco. (Plenary Abstract)
27. Schusterman, R.J. and Kastak, D. (1998) Recent advances in marine mammal acoustics: noise pollution, habitat degradation, and policy making. The World Marine Mammal Science Conference, Monaco. (Symposium Abstract)
28. Kastak, D. and Schusterman, R.J. (1998) Pinniped hearing: amphibious adaptations and the effects of noise. The World Marine Mammal Science Conference, Monaco. (Abstract)
29. Southall, B.L., Kastak, D., and Schusterman, R.J. (1998) Masked underwater hearing thresholds in at low frequencies for two pinniped species. The World Marine Mammal Science Conference, Monaco. (Abstract)
30. Levenson, D.H. and Schusterman, R.J. (1998) Dark adaptation and visual sensitivity in phocid and otariid pinnipeds. The World Marine Mammal Science Conference, Monaco. (Abstract)